

# ATM 265, Spring 2019

## Lecture 13

### Global Atmospheric Models from Around the World

May 8, 2019

Paul A. Ullrich (HH 251)  
[paullrich@ucdavis.edu](mailto:paullrich@ucdavis.edu)


# Why Multiple Models?

**Figure:** Relative error metrics for 20<sup>th</sup> century CMIP3 annual cycle climatology (1980..1999). A value of -0.2 means that the model has an error 20% smaller than the typical model error for that quantity.

**Source:** NASA

In general, the model mean is a significantly more skillful approximation to the real climate than any individual model result.

*Why do you think this is the case?*



# *Operational Global Atm. Models*



IPCC participating model

# ***Global Atmospheric Models***

## **Outline:**

- Ocean-Land-Atmosphere Model (OLAM)
- Unified Z-Grid Icosahedral Model (CSU/UZIM)
- UK Met Office Unified Model / ENDGame
- Non-hydrostatic Icosahedral Cloud-resolving Atmosphere Model (NICAM)
- NCAR / CESM / Community Atmosphere Model (CAM)
- NCAR / Model for Prediction Across Scales (MPAS)
- DOE / E3SM
- ECMWF / Integrated Forecasting System (IFS)
- GFDL / Finite-Volume Cubed (FV3)
- Environment and Climate Canada / Global Environmental Multiscale Model (GEM)

# ***Ocean-Land-Atmosphere Model (OLAM)***

## **Authors:**

Robert Walko (University of Miami, Miami, FL)

Roni Avissar (University of Miami, Miami, FL)

Martine Otte (US Environmental Protection Agency)


## **Operational Model, Non-Hydrostatic, Deep Atmosphere**

**Used by:** US Environmental Protection Agency

**Based on:** Limited-Area Model RAMS (1986)


# OLAM

The model is based on an icosahedral triangular or hexagonal grid.




# OLAM

Cut cell method used in the vertical (improves horizontal pressure gradient calculation)



Cut cells treated by scaling surface area of cells by “exposed” area.



# ***Colorado State University Unified Z-grid Icosahedral Model (UZIM)***

## **Authors:**

Dave Randall (Colorado State University, Fort Collins, CO)

Don Dazlich (Colorado State University, Fort Collins, CO)

Ross Heikes (Colorado State University, Fort Collins, CO)

## **Non-Operational Model, Non-Hydrostatic, Shallow Atmosphere**

## **Based on:**

Unified model system (Arakawa and Konor 2009)



Vorticity-Divergence Z-grid (Randall 1994)



The model is based a hexagonal grid which has been optimized via a post-processing step.

**Z-Grid formulation:**  
Each hexagon stores the vorticity and divergence at that point (instead of the zonal and meridional velocities).

This has been shown to have better numerical properties, but requires a global solve at each time step to recover  $u, v$ .



# ***UZIM***

- Third-order Multi-Step scheme (Adams Bashforth) for time stepping
- Positivity preserving Flux-Corrected Transport (FCT)
- “Unified” approach for treatment of vertically propagating sound waves
- Topography not yet implemented (plan is for cut cells)

# **ENDGame**

## **UK Met Office Model**

### **Authors:**

John Thuburn (University of Exeter)

Thomas Melvin (UK Met Office)

Many Others...

## **Operational Model, Non-Hydrostatic/Hydrostatic, Shallow/Deep Atmosphere**


**Philosophy:** “Unified” model used for both Numerical Weather Prediction applications and Climate applications. Must satisfy NWP benchmarks: Must forecast at 25km 7 days in < 3 hours.

# ENDGame



The model is based a latitude-longitude grid. The small timestep restriction is overcome using a semi-Lagrangian methodology.

- Includes modification to ellisoidal geopotential.
- C-Grid in the horizontal
- Charney-Phillips vertical staggering
- Fully implicit (iterative) scheme for non-advection terms
- Scheme not inherently conservative (a posteriori correction scheme needed)



# ***Nonhydrostatic Icosahedral Cloud-resolving Atmospheric Model (NICAM)***

**University of Tokyo Cloud Resolving Model**

**Authors:**

Masaki Satoh (University of Tokyo)

Hirofumi Tomita (Japan Agency for Marine-Earth Science and Technology)

Hiroaki Miura (University of Tokyo)

Ryuji Yoshida (AICS, Riken, Kobe)


Many Others...

**Operational Model, Non-Hydrostatic, Shallow/Deep Atmosphere**

**Philosophy:** Design a model for cloud resolving scales (3 km) global resolution.

(Note: Experimental runs have been produced at 800m)

# NICAM



Global triangular icosahedral mesh.



Madden-Julien Oscillation Experiment  
Left: Satellite, Right: NICAM.

- **Project goal:** Resolve horizontal scales well enough to not require a cumulus cloud parameterization.
- Model was among several designed to take advantage of the super-computing architecture of the Earth Simulator.
- **Highly local finite difference method:** Project aimed to use the simplest numerical method possible and maximize computational efficiency.
- A-grid finite-volume method with 2<sup>nd</sup> order central spatial discretizations plus terrain-following vertical coordinate. Split-explicit treatment of sound waves.

# ***Integrated Forecast System (IFS)***

**European Center for Medium-Range Weather Forecasting (ECMWF)**

**Authors:**

Nils Wedi (ECMWF)

Sylvie Malardel (ECMWF)

Many others...

**Operational Model, Hydrostatic/Non-hydrostatic, Shallow/Deep Atmosphere**

**Philosophy:** High-resolution high-throughput numerical weather prediction system using tried-and-tested numerical methods.

# *Integrated Forecast System (IFS)*




Spectral transform  
method based on a  
latitude-longitude grid.

Year	Truncation	Resolution
1983	T63	~ 316 km
1987	T106	~ 188 km
1991	T213	~ 95 km
1998	T <sub>L</sub> 319	~ 63 km
2000	T <sub>L</sub> 511	~ 39 km
2006	T <sub>L</sub> 799	~ 25 km
2010	T <sub>L</sub> 1279	~ 16 km
2015	T <sub>L</sub> 2047	~ 9 km*
2020	.. T <sub>L</sub> 3999	1 – 9 km**


\* Hydrostatic, parameterized convection

\*\* Non-hydrostatic, explicit convection

# *Integrated Forecast System (IFS)*



Northern hemisphere regular longitude-latitude grid.



Northern hemisphere reduced longitude-latitude grid (30% reduction in the number of grid points)

# **Community Earth System Model (CESM)**

# **Community Atmosphere Model (CAM)**

**National Center for Atmospheric Research (NCAR)**

**Authors:**

David Williamson, Rich Loft, Rich Neale, Peter Lauritzen, etc. (NCAR)

Many others...

**Operational Model, Hydrostatic, Shallow Atmosphere**

**Philosophy:** A community resource for studying the atmosphere. Focus is on climate applications. Large-scale parallel scalability for the CAM-SE model.

# *Energy Exascale Earth System Model (E3SM) / E3SM Atmosphere Model (EAM)*

**Department of Energy (DOE)**

**Authors:**

Dave Bader, Peter Caldwell, Mark Taylor, etc. (DOE)

Many others...

**Operational Model, Hydrostatic, Shallow Atmosphere**

**Philosophy:** A model branched from CESM targeted at extreme scale computing and answering questions related to DOE interests. Dynamical core extended to non-hydrostatic in 2019.

# Community Atmosphere Model (CAM)

Support for four dynamical cores:

- ***Spectral element (default)*** – Default in CAM 5.2
- ***Finite-Volume (previous default)*** – Used in IPCC AR5 and AR6
- ***Semi-Lagrangian (deprecated)***
- ***Global Spectral Transform (Eulerian dynamics)*** – Used in IPCC AR4



Spectral Element



Finite-Volume  
Semi-Lagrangian  
Eulerian

# Community Atmosphere Model (CAM)

In the vertical:

- Hybrid sigma-pressure coordinate
- Floating Lagrangian layers
- All physical parameterizations must work within a column (only column physics)





Figure: Nair et al. (2009)


# Community Atmosphere Model (CAM)

CAM-SE is built as a completely unstructured model on a conformal grid.

Using mesh refinement, essentially any grid shape is supported.



Arbitrarily shaped quadrilateral elements supported by spectral element method.



# *Model for Prediction Across Scales (MPAS)*

National Center for Atmospheric Research (NCAR)

Authors:

Bill Skamarock (NCAR)

Joe Klemp (NCAR)

Todd Ringler (Los Alamos National Laboratory)

Many others...

**Non-Operational Model, Non-Hydrostatic, Shallow/Deep Atmosphere**

**Philosophy:** Large-scale parallel scalability (next-generation atmospheric dynamics code) and support for variable resolution meshes.

# MPAS




The model is based a hexagonal grid which has been optimized via a post-processing step.

## C-Grid formulation:


Each hexagon stores the density and potential temperature at that point.

Velocities are stored as perpendicular components at each edge.

**Issues:** Number of velocity components is greater than number of density components.



# MPAS



**Figure:** An example of a variable-resolution centroidal Voronoi mesh used in MPAS.

# ***Finite-Volume Cubed (FV3)***

**Geophysical Fluid Dynamics Laboratory (GFDL)**

**Authors:**

Shian-Jiann (SJ) Lin (GFDL)

Richard Rood (University of Michigan)

Lucas Harris (GFDL)

Many others...

**Operational Model, Hydrostatic, Shallow Atmosphere**

**Experimental Non-hydrostatic mode**


**Other GFDL models:**

AM3/CM3, HiRAM, CM2.5/2.6, LASG, Academia Sinica, GISS ModelE


# FV3



Third-order finite-volume method on the cubed-sphere grid. Accuracy drops to second-order due to dimension splitting.



Arakawa D-grid  
staggering for velocities.



Floating Lagrangian layers

# ***Global Environmental Multiscale Model (GEM)***

**Environment Canada / Canadian Meteorological Center (CMC)**

**Authors:**

Abdessamad Qaddouri (Environment Canada)

Jean Côté (Environment Canada)


Sylvie Graval (Environment Canada)

Paul Vaillancourt (Environment Canada)

Many others...

**Operational Model, Hydrostatic/Non-hydrostatic, Shallow Atmosphere**

# GEM



## Modified Charney-Phillips (“CP”) grid

